

Combinatorial Chemistry Online

Volume 9, Issue 1, January 2007

N. K. Terrett

Ensemble Discovery Corp., Cambridge, MA 02139, USA

1. Current literature highlights

1.1. Pyrazole CCK₁ receptor antagonists


Cholecystokinin (CCK) is an endogenous 33 amino acid peptide hormone first identified by its pharmacological actions early in the last century, and later purified and sequenced in the 1970s. Subsequently, it was shown that CCK was released in response to food intake and that it regulates gallbladder contraction, pancreatic enzyme secretion, gastric acid secretion, and colonic motility amongst other pharmacological effects. CCK is abundant in the CNS and is believed to be involved in aspects of nociception, satiety and anxiogenesis.

The biological actions of CCK are mediated through two G-protein coupled receptors, CCK₁ and CCK₂, and a number of CCK₁ antagonists have been evaluated in the clinic for pancreatic disorders, IBS and biliary colic. Encouraging clinical results from a phase II trial of constipation-dependent IBS with a peptide derived CCK₁ antagonist has given encouragement to pursue non-peptide antagonists of CCK₁.¹

In this recent work, efforts were undertaken to modify an HTS lead (i) that was identified as a potent and selective antagonist of the CCK₁ receptor (pK_i 7.6). This compound displayed promising physical properties and represented a novel chemical scaffold for SAR investigations undertaken by the construction of a solution phase library. The synthetic scheme involved varying positions around aromatic centers B and C giving antagonists of CCK₁ with pK_i values between 6.2 and 8.1.

During the course of these studies, a novel method for the evaluation of SAR in combinatorial matrices was explored. This new methodology provided for a quantitative assessment of additive and non-additive relationships in the

SAR, allowing one to identify potential changes in the binding modes of these antagonists. The assumption with additive SAR models is that one variable in the structure does not affect the binding or conformation of the second variable. However, in many documented cases this assumption is invalid, thus limiting the predictive power of additive models. With the advent of combinatorial methods, a full matrix of compounds can be readily accessed synthetically, and the presence of non-additive relationships in the full matrix can suggest either a different binding mode in the biological target or direct interactions between different parts of a ligand. The results from this study indicate that all compounds tested bind to the receptor in a similar binding mode.



1.2. Aminopyrrolidinetricarboxylic acids: new group III metabotropic glutamate receptor-selective antagonists

The glutamate neurotransmitter plays a critical role in many disease processes. Glutamate exerts its effects through two families of receptors: metabotropic (mGlu) and ionotropic (iGlu) receptors. mGlu receptors have been further divided in three groups according to their sequence homologies, pharmacological properties and signal transduction pathways.


Group III mGlu receptors include mGlu4, mGlu6, mGlu7 and mGlu8 receptors, which are negatively coupled to cAMP production. So far, only a few orthosteric ligands, that compete with glutamate, have been described for

E-mail: nterrett@ensemlediscovery.com

group III mGlu receptors. Typically, these agonists have low-micromolar to sub-micromolar potencies for mGlu4, mGlu6 and mGlu8, whilst displaying much less potency at the mGlu7 receptor. Recent work has demonstrated the utility of parallel chemistry for the rapid synthesis of polar amino acid derivatives that are orthosteric agonists of group III mGlu receptors.²

This work focuses on aminopyrrolidine-tricarboxylic acid (APTC) derivatives based on a family of APTCs known to provide selective group III agonists. The aim was to have a rigid template which would allow for the placement of polar functionality around the core, thus encouraging interaction of these polar groups with distal binding pockets in the receptors. Following solution phase synthesis of diastereoisomerically-pure protected cores, exemplified by generic structure (ii), a library of 65 compounds was constructed in solution resulting from capping of the pyrrolidine nitrogen with various (R)-groups, followed by deprotection of the Boc protecting-group. The nitrogen was capped with electrophilic reagents using solid supported reagents and scavengers.

The compounds prepared were screened against three group III mGlu rat receptors (mGlu4, mGlu6 and mGlu8) for their agonist and antagonist activities at 450 μ M and 300 μ M, respectively. One of the most potent compounds obtained was (iii) which displayed an agonist response to mGlu4 with an EC_{50} of 48 μ M. This compound displayed 7-fold selectivity against mGlu6 and was equally active against mGlu8. Compound (iii) was inactive ($EC_{50} > 5000$ μ M) against group I receptors (mGlu1, mGlu5) and the group II receptor mGlu2. Interestingly, compound (iii) is a full agonist for mGlu4 but a partial agonist for mGlu8, despite it having a similar EC_{50} value against both mGlu4 and mGlu8.



2. A summary of the papers in this month's issue

2.1. Solid-phase synthesis

A general and efficient solid-phase synthesis of N-9-substituted 2,8-diaminopurines from 5-nitrouracil has been described. The key synthetic transformation employs a carbodiimide-mediated cyclisation of a thiourea.³

The solid-phase synthesis of fused [2,1-*b*]quinazolinone alkaloids has been developed for the preparation of vasicinone and deoxyvasicinone by two alternative approaches.⁴

A new method for the preparation of peptide thioester by the post-solid phase peptide synthesis (SPPS) approach

has been developed. A series of *N*-alkyl cysteine derivatives were prepared and used as the C-terminus residue of the peptides prepared by the Fmoc SPPS, and the synthetic peptides released from resin by TFA were readily converted to the peptide thioester without significant side reactions.⁵

A method to transform allyl esters to thioesters under solid phase conditions has been developed to synthesise peptide thioesters.⁶

A microwave-assisted traceless rapid synthesis of benzimidazolones on polymeric supports has been developed.⁷

Two novel microgonotropens (MGTs) comprised of hairpin *N*-propylaminepyrrole polyamides linked to a Hoechst 33258 (Ht) analogue have been synthesised on solid phase by adopting an Fmoc technique using a series of HOBT mediated coupling reactions.⁸

2.2. Solution-phase synthesis

Di-*exo*-3-amino-7-oxabicyclo[2.2.1]hept-5-ene-2-carboxylic acid, five aldehydes and two isocyanides were reacted both in methanol and in water to prepare a 10-membered β -lactam library via an Ugi-4-centre-3-component reaction.⁹

The concept, protocols and instruments of fluorous solid-phase extraction techniques, and their applications for solution-phase parallel and high-throughput synthesis of small molecules and biomolecules have been reviewed.¹⁰

Nucleophilic addition of Grignard or lithiated reagents to new Weinreb amides efficiently gave the corresponding ketones and allowed the design and synthesis of a new indazole library.¹¹

A rapid library-generation via liquid-phase multiple-parallel synthesis of 2-(substituted)benzyl-1-benzopyrano[4,3-*c*]pyrazol-3(2*H*)-ones, bearing two points of diversity, under microwave irradiation has been successfully performed using chromenone-3-carboxylic acids as starting materials. Compared to an identical library generated by conventional parallel synthesis, microwave-assisted parallel synthesis dramatically decreased reaction times from an average of 16 h to 13 min, and the yields of products and intermediates were improved in most cases.¹²

2.3. Scaffolds for combinatorial libraries

The synthesis of a 5,7-dichloropyrido[4,3-*d*]pyrimidine scaffold has been described. The chlorine at position 5 can selectively be displaced by different palladium-catalyzed cross-coupling reactions and nucleophilic aromatic substitutions. In the subsequent step, the chlorine at position 7 can be further derivatised, allowing for the construction of a diverse pyrido[4,3-*d*]pyrimidine library with structural variations at positions 5 and 7.¹³

2.4. Solid-phase supported reagents

The application of a 'catch and release' approach to palladium-catalysed multi-component cascade reactions leads to

diverse libraries of pharmacologically interesting small molecules in high yield and with excellent purity.¹⁴

Polymer-supported methyltrioxorhenium (MTO) systems are efficient catalysts for the oxidative functionalisation of cyclohexane and cyclopentane derivatives with H₂O₂ as oxygen donor.¹⁵

2.5. Novel resins, linkers and techniques

A solid macroporous monolith is shown to be a suitable substrate for anchoring a palladium complex to obtain a continuous porous material suitable for conducting flow-through catalysis in capillary microreactors.¹⁶

Straightforward syntheses of two *tert*-alkoxysilyl chloride functionalised resins that allow facile attachment of 1°, 2°, 3° alcohols and phenols to the solid-phase have been achieved.¹⁷

2.6. Library applications

A novel bacterial transcription/translation (TT) inhibitor has been identified through a combination of high throughput screening and exploratory medicinal chemistry. Initial optimisation of an anthranilic acid moiety and sulphonamide amine diversity was accomplished via one- and two-dimensional solution phase libraries.¹⁸

Solid-phase synthesis of a small combinatorial library of dihydroceramide analogues as mixtures of *erythro* and *threo* diastereomers has been described. Some dihydroceramide analogues cause growth arrest and apoptosis in a dose-dependent manner in human alveolar epithelial cells.¹⁹

The design and parallel synthesis of 217 compounds based on a malonic-hydroxamic acid template has been reported. These compounds are obtained via a two-step solution-phase procedure, and the diverse building-blocks used makes this strategy suitable for the search for inhibitors of various metallo-proteases and for the investigation of the biological role of new metallo-proteases.²⁰

T-type calcium channel is one of therapeutic targets for the treatment of cardiovascular diseases and neuropathic pains. A small molecule library of dioxoquinazoline carboxamide derivatives containing 155 compounds was designed, synthesised, and biologically evaluated for T-type calcium channel blocking activity.²¹

Cu(I)-catalysed ‘click’ cycloaddition reactions between azides and alkynes have been employed to generate two sequential libraries of protein tyrosine phosphatase (PTP) inhibitors. Products were screened against the *Yersinia* PTP and PTP1B, and four compounds were selected and evaluated in pure form against the *Yersinia* PTP, PTP1B, TCPTP, LAR and CD45.²²

A small library with more than forty diverse analogues of Oenostacin, a naturally occurring potent antibacterial phenolic acid derivative, have been prepared through solution

phase synthesis. Some of these analogues possessed potent antibacterial activities against *Staphylococcus epidermidis* and *Staphylococcus aureus*.²³

References

1. McClure, K. et al. *Bioorg. Med. Chem. Lett.* **2006**, 16 (1), 72–76.
2. Schann, S. et al. *Bioorg. Med. Chem. Lett.* **2006**, 16 (18), 4856–4860.
3. Cole, A. G. et al. *Tetrahedron Lett.* **2006**, 47 (50), 8897–8900.
4. Kamal, A. et al. *Tetrahedron Lett.* **2006**, 47 (51), 9025–9028.
5. Hojo, H. et al. *Tetrahedron Lett.* **2007**, 48 (1), 25–28.
6. Li, L.; Wang, P. *Tetrahedron Lett.* **2007**, 48 (1), 29–32.
7. Xu, X.-J.; Zong, Y.-X. *Tetrahedron Lett.* **2007**, 48 (1), 129–132.
8. Kahane, A. L.; Bruice, T. C. *Bioorg. Med. Chem. Lett.* **2006**, 16 (24), 6255–6261.
9. Kanizsai, I. et al. *Tetrahedron Lett.* **2006**, 47 (51), 9113–9116.
10. Zhang, W.; Curran, D. P. *Tetrahedron* **2006**, 62 (51), 11837–11865.
11. Crestey, F. et al. *Tetrahedron* **2007**, 63 (2), 419–428.
12. Zhou, Z.-Z.; Yang, G.-F., et al. *Bioorg. Med. Chem.* **2006**, 14 (24), 8666–8674.
13. Jang, M.-Y. et al. *Tetrahedron Lett.* **2006**, 47 (50), 8917–8920.
14. Grigg, R.; Cook, A. *Tetrahedron* **2006**, 62 (52), 12172–12181.
15. Bianchini, G. et al. *Tetrahedron* **2006**, 62 (52), 12326–12333.
16. Bolton, K. F. et al. *Tetrahedron Lett.* **2006**, 47 (52), 9321–9324.
17. Meloni, M. M. et al. *Tetrahedron* **2007**, 63 (2), 299–311.
18. Larsen, S. D. et al. *Bioorg. Med. Chem. Lett.* **2006**, 16 (24), 6173–6177.
19. Villorquina, G. et al. *Bioorg. Med. Chem.* **2007**, 15 (1), 50–62.
20. Flipo, M. et al. *Bioorg. Med. Chem.* **2007**, 15 (1), 63–76.
21. Jo, M. N. et al. *Bioorg. Med. Chem.* **2007**, 15 (1), 365–373.
22. Xie, J.; Seto, C. T. *Bioorg. Med. Chem.* **2007**, 15 (1), 458–473.
23. Srivastava, V. et al. *Bioorg. Med. Chem.* **2007**, 15 (1), 518–525.

Further reading

Papers on combinatorial chemistry or solid-phase synthesis from other journals

- Sun, L.-P.; Dai, W.-M. An engineered linker capable of promoting on-resin reactions for microwave-assisted solid-phase organic synthesis. *Angewandte Chemie, International Edition* **2006**, 45 (43), 7255–7258.
- Diness, F.; Beyer, J.; Meldal, M. Solid-phase synthesis of tetrahydro- β -carboline and tetrahydroisoquinolines by stereoselective intramolecular N-carbamyliminium Pictet-Spengler reactions. *Chemistry—A European Journal* **2006**, 12 (31), 8056–8066.
- Lesma, G.; Danieli, B.; Lodroni, F.; Passarella, D.; Sacchetti, A.; Silvani, A. Microwave-assisted, solid-phase synthesis of a chiral 1,2,3,4-tetrahydroquinoline library. *Combinatorial Chemistry & High Throughput Screening* **2006**, 9 (9), 691–701.
- Tye, H.; Whittaker, M.; Ramdeehul, S. Minimisation of palladium content in Suzuki cross-coupling reactions. *Combinatorial Chemistry & High Throughput Screening* **2006**, 9 (9), 703–710.
- Risseeuw, M. D. P.; Grotenbreg, G. M.; Witte, M. D.; Tuin, A. W.; Leeuwenburgh, M. A.; Van der Marel, G. A.; Overkleeft,

- H. S.; Overhand, M. The synthesis of cis- and trans-fused bicyclic sugar amino acids. *European Journal of Organic Chemistry* **2006**, (17), 3877–3886.
- Gea, A.; Farcy, N.; Roque i Rossell, M.; Martins, J. C.; De Clercq, P. J.; Madder, A. Solid-supported synthesis of highly functionalized tripodal peptides with flexible but preorganized geometry: towards potential serine protease mimics. *European Journal of Organic Chemistry* **2006**, (18), 4135–4146.
- Joo, S. H.; Xiao, Q.; Ling, Y.; Gopishetty, B.; Pei, D. High-throughput sequence determination of cyclic peptide library members by partial Edman degradation/mass spectrometry. *Journal of the American Chemical Society* **2006**, 128 (39), 13000–13009.
- Le Lamer, A.-C.; Gouault, N.; David, M.; Boustie, J.; Uriac, P. Method for the parallel synthesis of α -methylene- γ -lactones from a fluorous acrylate. *Journal of Combinatorial Chemistry* **2006**, 8 (5), 643–645.
- Sharma, U.; Srivastava, S.; Prakesch, M.; Sharma, M.; Leek, D. M.; Arya, P. Part 2: Building diverse natural product-like architectures from a tetrahydroaminoquinoline scaffold. Modular Solution- and Solid-Phase Approaches for Use in High-Throughput Generation of Chemical probes. *Journal of Combinatorial Chemistry* **2006**, 8 (5), 735–761.
- Prakesch, M.; Srivastava, S.; Leek, D. M.; Arya, P. Part 3. A novel stereocontrolled, in situ, solution- and solid-phase, aza Michael approach for high-throughput generation of tetrahydroaminoquinoline-derived natural product-like architectures. *Journal of Combinatorial Chemistry* **2006**, 8 (5), 762–773.
- Kowalski, J. A.; Leonard, S. F.; Lee, G. E., Jr. Diverse 2-carboxamide-3-amino-substituted quinoxalines: synthesis and reactivity investigation for library generation. *Journal of Combinatorial Chemistry* **2006**, 8 (5), 774–779.
- Jeon, H. S.; Kim, J. N.; Kim, T. H. Solid-phase synthesis of 2-amino-2-thiazolines from N-(2-hydroxyethyl)thioureas using 1,3-dicyclohexylcarbodiimide. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 799–801.
- Sheng, S.-R.; Xu, L.; Zhang, X.-L.; Liu, X.-L.; Wei, M.-H. Solid-phase synthesis of substituted butenolides and butyrolactones using a traceless sulfone linker. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 805–807.
- Cardullo, F.; Donati, D.; Fusillo, V.; Merlo, G.; Paio, A.; Salaris, M.; Solinas, A.; Taddei, M. Parallel protocol for the selective methylation and alkylation of primary amines. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 834–840.
- Kou, B.-B.; Zhang, F.; Yang, T.-M.; Liu, G. Simultaneous solid-phase synthesis of quinoxalinone and benzimidazole scaffold libraries. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 841–847.
- Reddy, P. T.; Quevillon, S.; Gan, Z.; Forbes, N.; Leek, D. M.; Arya, P. Solution- and solid-phase, modular approaches for obtaining different natural product-like polycyclic architectures from an aminoindoline scaffold for combinatorial chemistry. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 856–871.
- Doemling, A.; Beck, B.; Fuchs, T.; Yazbak, A. Parallel synthesis of arrays of amino-acid-derived isocyanoamides useful as starting materials in IMCR. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 872–880.
- Zhang, W.; Lu, Y. Automation of fluorous solid-phase extraction for parallel synthesis. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 890–896.
- Hwang, J. Y.; Choi, H.-S.; Seo, J.-s.; La, H.-J.; Yoo, S.-e.; Gong, Y.-D. Method for the solid-phase parallel synthesis of a 6-alkylamino-2-(functionalized-aminomethyl)-2H-1-benzopyran library. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 897–906.
- Tullberg, M.; Luthman, K.; Grotli, M. Microwave-assisted solid-phase synthesis of 2,5-diketopiperazines: solvent and resin dependence. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 915–922.
- He, R.; Ching, S. M.; Lam, Y. Traceless solid-phase synthesis of substituted xanthenes. *Journal of Combinatorial Chemistry* **2006**, 8 (6), 923–928.
- Kumar, A.; Ye, G.; Ahmadibeni, Y.; Parang, K. Synthesis of polymer-bound 4-acetoxy-3 phenylbenzaldehyde derivatives: applications in solid-phase organic synthesis. *Journal of Organic Chemistry* **2006**, 71 (20), 7915–7918.
- Bose, D. S.; Idrees, M. Hypervalent iodine mediated intramolecular cyclization of thioformanilides: expeditious approach to 2-substituted benzothiazoles. *Journal of Organic Chemistry* **2006**, 71 (21), 8261–8263.
- Testero, S. A.; Mata, E. G. Synthesis of 3-(aryl)alkenyl- β -lactams by an efficient application of olefin cross-metathesis on solid support. *Organic Letters* **2006**, 8 (21), 4783–4786.
- Fernandez, J.-C.; Sole-Feu, L.; Fernandez-Forner, D.; de la Figuera, N.; Forns, P.; Albericio, F. Solid-phase N-electrophilic amination of 2-aminopyridines: preparation of 2-substituted-[1,2,4]triazolo[1,5-a]pyridine derivatives. *QSAR & Combinatorial Science* **2006**, 25 (10), 961–965.
- Monroc, S.; Feliu, L.; Serra, J.; Planas, M.; Bardaji, E. Synthesis and solid-phase applications of N-tetrachlorophthaloyl (TCP) side-chain-protected amino acids. *Synlett* **2006**, (17), 2743–2746.
- Furdui, B.; Dinica, R.; Druta, I. I.; Demeunynck, M. Improved synthesis of cationic pyridinium-substituted indolizines. *Synthesis* **2006**, (16), 2640–2642.
- Zinieris, N.; Kokinaki, S.; Leondiadis, L.; Ferderigos, N. Synthesis of [4-(Fmoc-aminoacyloxymethyl)phenoxy]acetic acids for use in solid-phase peptide synthesis. *Synthesis* **2006**, (16), 2789–2793.
- Boggian, D. B.; Mata, E. G. A versatile strategy for the solid-phase synthesis of penicillin derivatives: efficient preparation of 2 β -methyl substituted penams as β -lactamase inhibitor analogues. *Synthesis* **2006**, (20), 3397–3404.